



O'ZMU XABARLARI

ВЕСТНИК НУУЗ

АСТА NUUZ

MIRZO ULUG'BEK NOMIDAGI O'ZBEKISTON MILLIY
UNIVERSITETINING ILMIIY JURNALI

**JURNAL 1997
YILDAN
CHIQA
BOSHLAGAN**

**2025
2/1/1
Aniq
fanlar**

Bosh muharrir:

MADJIDOV I. U. – t.f.d., professor

Bosh muharrir o'rinbosari:

ERGASHOV Y. S. – f.-m.f.d., professor

Tahrir hay'ati:

- Ayupov Sh. A.** – f.-m.f.d., prof., O'zR FA akademigi
Alimov Sh. A. – f.-m.f.d., prof., O'zR FA akademigi
Sadullayev A. – f.-m.f.d., prof., O'zR FA akademigi
Roziqov O'. A. – f.-m.f.d., prof., O'zR FA akademigi
Aripov M. M. – f.-m.f.d., prof.
Zikirov O. S. – f.-m.f.d., prof.
Abdushukurov A.A. – f.-m.f.d., prof.
Aloyev R. J. – f.-m.f.d., prof.
Ganixodjayev R. N. – f.-m.f.d., prof.
Narmonov A. Y. – f.-m.f.d., prof.
Raximov A. A. – f.-m.f.d., prof.
Beshimov R. B. – f.-m.f.d., prof.
Shoimqulov B. A. – f.-m.f.d., prof.
Axmedov A.B. – f.-m.f.d. prof.
Sharipov O.Sh. – f.-m.f.d., prof.
Mamadaliyev N.A. – f.-m.f.d., prof.
Xudoyberdiyev A.X. – f.-m.f.d., prof.
Xudoyberganov G. – f.-m.f.d., prof.
Matyakubov A. S. – f.-m.f.d., prof.

Mas'ul kotib: f.-m.f.d. (PhD) dots. **G'aybullayev R.Q.**

TOSHKENT – 2025

◇ MUNDARIJA ◇ CONTENTS ◇ СОДЕРЖАНИЕ ◇

Abdukaxorova Z. T. H_{A_2} -weakly periodic p -adic generalized gibbs measures for the ising model with an external field on the cayley tree of order two	3
Akhmedova Q., Yusupova Sh., Mirzoodilov J. Qashqadaryo turizmining turg'un rivojlanishida graflar nazariyasining qo'llanilishi	13
Buvayev Q. T., Davirova G. N. Furrye qatorlarining elliptik qisman yig'indilar ketma-ketligi uchun umumlashgan lokalizatsiya masalasi	19
Dusmatov O. M., Xodjabekov M. U., Kasimova F. U. Nomukammal elastik sterjenning chiziqalmas tebranishlari ustuvorligi	31
Gaybullaev R. K., Solijanov G. O., Usmonov J. B. 2D moore ca with new boundary conditions and its reversibility	39
Ibragimov H. H. Pifagor uchligining fraktal xossalari	50
Khusanbaev Ya. M., Toshkulov X. A. On the asymptotic of branching reduced processes ...	59
Lakaev Sh. S., Aktamova V. U. On the point spectrum of the Schrodinger operator for a system consisting of two identical infinitely heavy bosons and one light fermion on the three-dimensional lattice	66
Matanov M. Ch. Harmonic fundamental solutions and boundary integral equations for wave propagation in elastic, scalar and poroelastic media	80
Mukhamadiev F. G., Meyliev Sh. U. Some homotopy properties of n -fold symmetric product of the space X	88
Normatov E. P., Elboyev K. E. \mathbb{Z}_2 maydonda ikki va uch o'lchamli evolyutsion algebralarning tasnifi	92
Ruzieva D. S. Strong law of large numbers for Banach space-valued random fields with weights	106
Safarov U. A. Some remark on the invariant measure of circle maps with break points	111
Sharipov O. Sh., Kobilov U. X. On stable type p Banach spaces	115
Sharipov O. Sh., Kushmurodov A. A. Strong law of large numbers for dependent random variables with values in Rademacher type p Banach spaces	120
Zhabborov N. M., Tuychiyev S. G., Eshdavlatova S. E. Estimating the unemployment rate using mathematical modeling	125
Ахмедов А. Б., Ибрагимова Н.И. Теория равновесия пластин в напряжениях	132
Давлатбеков А. А., Собирова М. Р. О нормальных формах и некоторых свойствах линейных квазигрупп и квазигрупп смешанного типа первого (второго) рода	142
Зупаров Т.М., Жовлиев А.И. Центральная предельная теорема и оценка остаточного члена в центральной предельной теореме (ц.п.т) для линейных процессов	147
Касимов Ш. Г., Коцанов А. П. Об однозначной разрешимости многомерной начально-граничной задачи для уравнения высокого порядка с дробной производной в смысле Миллера - Росса в классах Соболева	155
Мадрахимов Р. М., Омонов О. И. Свойства M -субгармонических функций	165
Мамадалиев Н. А., Васиева Х. Г. Управление пучками траекторий в квазилинейных дифференциально-разностных играх нейтрального типа	170
Мустапокулов Х. Я. Задачи преследования-убегания для дифференциальных игр второго порядка с импульсным управлением	178
Рахимов А. М., Имомназаров Б. Х., Искандаров И. К. Задача коши для динамической системы пороупругости описываемой неоднородными уравнениями с тремя параметрами упругости	185
Рахмонов Ф. Д., Холтураев Ф. С. Смешанная задача для интегро-дифференциального уравнения типа Бенни-Люк высокого порядка с вырожденным ядром	191
Рахмонов Ш. М. Геометрия многообразия Грассмана	200
Садуллаев А., Шарипов Р. Оценка гессианов ограниченных m -выпуклых функций	206
Файзиев А. А. Статистический анализ и прогнозирование динамики заготовленного хлопка Сирдарьинской области в Республике Узбекистан	213

UDC 517.55

SOME HOMOTOPY PROPERTIES OF n -FOLD SYMMETRIC PRODUCT OF THE SPACE X

MUKHAMADIEV F. G.

 NATIONAL UNIVERSITY OF UZBEKISTAN, UZBEKISTAN NATIONAL PEDAGOGICAL UNIVERSITY, TASHKENT
 farhodgm@nuu.uz

MEYLIEV SH. U.

 NATIONAL UNIVERSITY OF UZBEKISTAN, TASHKENT
 shmeyliev@mail.ru

RESUME

In this paper, we study some homotopy properties of the space n -fold symmetric product of the space X . We prove that if mappings $f, g : X \rightarrow Y$ are homotopic, then the mappings $\mathcal{F}_n f, \mathcal{F}_n g : \mathcal{F}_n X \rightarrow \mathcal{F}_n Y$ are also homotopic. Also shown that the functor of n -fold symmetric product \mathcal{F}_n is a covariant homotopy functor. Besides proved that if spaces X and Y are homotopically equivalent, then the spaces $\mathcal{F}_n X$ and $\mathcal{F}_n Y$ are also homotopically equivalent.

Key words: Functor, n -fold symmetric product, homotopy, homotopically equivalent.

Recently, the topological properties on hyperspaces with the Vietoris topology and the homotopy properties of the topological spaces have been studied by many authors ([1], [2], [3], [4], [5], [6]).

In [1] the connection between a finally compact, pseudocompact, extremely disconnected, \aleph -space and its hyperspace is studied. And in the work [2] have been studied the connection between a uniformly connected, uniformly pseudocompact, P -precompact and its hyperspace. In the works [3] and [4] have been studied some cardinal and homotopy properties of the superextension λX of a topological space X . And in [4] proved that the superextension functor λ preserves homotopy, i.e. that it is a homotopy functor. In [5] showed that the functor of Permutation Degree SP_G^n preserves the homotopy and the retraction of topological spaces. And in [6] have been studied some homotopy properties of the space of complete linked systems.

Recall that a covariant functor is a mapping \mathcal{F} which assigns to a topological space X the space $\mathcal{F}(X)$, and to a continuous mapping $f : X \rightarrow Y$, the mapping $\mathcal{F}(f) : \mathcal{F}(X) \rightarrow \mathcal{F}(Y)$ satisfying the following conditions:

- 1) \mathcal{F} preserves identity, that is, if id_X is the identity mapping of X , then $\mathcal{F}(id_X) = id_{\mathcal{F}(X)}$;
- 2) \mathcal{F} preserves composition, that is, if $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are continuous mappings, then we have

$$\mathcal{F}(g \circ f) = \mathcal{F}(g) \circ \mathcal{F}(f).$$

We refer the reader to the book [7] and the article [8] for more information about functors. Some metric properties of n -fold symmetric product of the space X is studied in the work [9]. In this paper we study some homotopy properties and retractions of n -fold symmetric product of the space X .

All of our space are Hausdorff unless otherwise indicated. The symbol N stands for the set of positive integers and R stands for the set of real numbers. Given a space X , we define its hyperspaces as the following sets:

- 1) $CL(X) = \{A \subset X \mid A \text{ is closed and nonempty}\}$;
- 2) $2^X = \{A \in CL(X) \mid A \text{ is compact}\}$;
- 3) $\mathcal{F}_n(X) = \{A \in 2^X \mid A \text{ has at most } n \text{ points}\}$, $n \in N$ (see [9, 10]).

$CL(X)$ is topologized by the Vietoris topology defined as the topology generated by

$$\beta = \{\langle U_1, \dots, U_k \rangle \mid U_1, \dots, U_k \text{ are open subsets of } X, k \in N\},$$

where $\langle U_1, \dots, U_k \rangle = \{A \in CL(X) \mid A \subset \bigcup U_j \text{ and } A \cap U_j \neq \emptyset \text{ for each } j \in \{1, \dots, k\}\}$.

Note that, by definition, 2^X , $\mathcal{F}_n(X)$ and $\mathcal{F}(X)$ are subsets of $CL(X)$. Hence, they are topologized with the appropriate restriction of the Vietoris topology. Moreover,

- 1) $CL(X)$ is called the *hyperspace of nonempty closed subsets of X* ;
- 2) 2^X is called the *hyperspace of nonempty compact subsets of X* ;
- 3) $\mathcal{F}_n(X)$ is called the *n -fold symmetric product of X* ;
- 4) $\mathcal{F}(X)$ is called the *hyperspace of finite subsets of X* .

On the other hand, it is obvious that $\mathcal{F}(X) = \bigcup_{n=1}^{\infty} \mathcal{F}_n(X)$ and $\mathcal{F}_n(X) \subset \mathcal{F}_{n+1}(X)$ for each $n \in \mathbb{N}$ (see [9, 10]).

Remark 1. Let X be a space and let $n \in \mathbb{N}$.

- 1) $\mathcal{F}_n(X)$ is closed in $\mathcal{F}(X)$;
- 2) $f_1 : X \rightarrow \mathcal{F}_1(X)$, $(x \mapsto x)$, is a homeomorphism;
- 3) Every $\mathcal{F}_m(X)$ is a closed subset of $\mathcal{F}_n(X)$ for each $m, n \in \mathbb{N}$, $m < n$ (see [11]).

Notation 1. If U_1, U_2, \dots, U_n are open subsets of a space X , then $\langle U_1, U_2, \dots, U_n \rangle_{\mathcal{F}(X)}$ denotes the intersection of the open set $\langle U_1, U_2, \dots, U_n \rangle$ of the Vietoris topology, with $\mathcal{F}(X)$ (see [12]).

Notation 2. Let X be a space. If $\{x_1, x_2, \dots, x_r\}$ is a point of $\mathcal{F}(X)$ and $\{x_1, x_2, \dots, x_r \in \langle U_1, U_2, \dots, U_n \rangle_{\mathcal{F}(X)}\}$, then for each $j \leq r$, we let $U_{x_j} = \cap \{U \in \{U_1, U_2, \dots, U_s\} : x_j \in U\}$. Observe that $\langle U_{x_1}, U_{x_2}, \dots, U_{x_r} \rangle_{\mathcal{F}(X)} \subset \langle U_1, U_2, \dots, U_s \rangle_{\mathcal{F}(X)}$ (see [13]).

For some undefined or related concepts, we refer the reader to [14], [15] and [16].

Now we will consider some homotopic properties of n -fold symmetric product of the space X . We begin with definitions of notions that will be used in this section. We mainly follow terminology from [15] and [16].

Continuous mappings $f, g : X \rightarrow Y$ are said to be *homotopic* if there is a continuous mapping $H : X \times I \rightarrow Y$ such that $H(x, 0) = f(x)$ and $H(x, 1) = g(x)$. The mapping H is called a *homotopy* between f and g and we write $f \simeq g$ [15].

Let $f : X \rightarrow Y$ is a continuous mapping and $A \in \mathcal{F}_n X$. Let's assume that $(\mathcal{F}_n f)(A) = f(A)$. Then the mapping $\mathcal{F}_n f : \mathcal{F}_n X \rightarrow \mathcal{F}_n Y$ also is a continuous mapping.

For the functor of n -fold symmetric product \mathcal{F}_n the following theorem holds.

Theorem 1. If mappings $f, g : X \rightarrow Y$ are homotopic, then the mappings $\mathcal{F}_n f, \mathcal{F}_n g : \mathcal{F}_n X \rightarrow \mathcal{F}_n Y$ are also homotopic.

Proof. Assume that the mappings $f, g : X \rightarrow Y$ are homotopic. Then there exists a continuous mapping $H : X \times I \rightarrow Y$ such that $H(x, 0) = f(x)$, and $H(x, 1) = g(x)$. On the other hand, we have that $(\mathcal{F}_n f)(A) = f(A)$ and $(\mathcal{F}_n g)(A) = g(A)$ for all $A \in \mathcal{F}_n X$. Now we can define the mapping $(\mathcal{F}_n H)(A, t) = H(a, t)$, where $A \in \mathcal{F}_n X$ and $t \in [0, 1]$. It is clear that since the mapping H is continuous, the mapping $\mathcal{F}_n H$ is also continuous. Now we will show that the mapping $\mathcal{F}_n H$ is a homotopy between the mappings $\mathcal{F}_n f$ and $\mathcal{F}_n g$. Indeed $(\mathcal{F}_n H)(A, 0) = H(A, 0) = f(A) = (\mathcal{F}_n f)(A)$, and $(\mathcal{F}_n H)(A, 1) = H(A, 1) = g(A) = (\mathcal{F}_n g)(A)$. This means that $\mathcal{F}_n f \simeq \mathcal{F}_n g$. Theorem 1 is proved.

From the Theorem 1 we get the following corollary.

Corollary 1. If the mapping H is homotopy between f and g , then the mapping also $\mathcal{F}_n H$ is a homotopy between $\mathcal{F}_n f$ and $\mathcal{F}_n g$.

A continuous mapping $f : X \rightarrow Y$ is said to be a *homotopy equivalence* if there exists a continuous mapping $g : Y \rightarrow X$ such that the compositions $g \circ f$ and $f \circ g$ are homotopic to the identity mappings on X and Y respectively. Two topological spaces X and Y is said to be *homotopically equivalent* (notation, $X \simeq Y$) if there exists a homotopy equivalence $f : X \rightarrow Y$ (see [15]).

Proposition 1. If the mapping $f : X \rightarrow Y$ is homotopy equivalence, then the mapping $\mathcal{F}_n f : \mathcal{F}_n X \rightarrow \mathcal{F}_n Y$ is also homotopy equivalence.

Proof. Suppose that the spaces X and Y are homotopically equivalent. Then there exists two continuous mappings $f : X \rightarrow Y$ and $g : Y \rightarrow X$ such that $f \circ g \simeq id_Y$ and $g \circ f \simeq id_X$. It means that there are two homotopy $H_1(y, t)$ and $H_2(x, t)$ such that $H_1(y, 0) = (f \circ g)(y)$, $H_1(y, 1) = y$, and $H_2(x, 0) = (g \circ f)(x)$, $H_2(x, 1) = x$. We can define the compositions $\mathcal{F}_n f \circ \mathcal{F}_n g : \mathcal{F}_n Y \rightarrow \mathcal{F}_n Y$, and $\mathcal{F}_n g \circ \mathcal{F}_n f : \mathcal{F}_n X \rightarrow \mathcal{F}_n X$ of the mappings $\mathcal{F}_n f : \mathcal{F}_n X \rightarrow \mathcal{F}_n Y$ and $\mathcal{F}_n g : \mathcal{F}_n Y \rightarrow \mathcal{F}_n X$ as follows: $(\mathcal{F}_n f \circ \mathcal{F}_n g)(B) = (f \circ g)(B)$ for all $B \in \mathcal{F}_n Y$, and $(\mathcal{F}_n g \circ \mathcal{F}_n f)(A) = (g \circ f)(A)$ for all $A \in \mathcal{F}_n X$. One can easily check that the mapping $(\mathcal{F}_n H_1)(B, t) = H_1(B, t)$ is a homotopy between $\mathcal{F}_n f \circ \mathcal{F}_n g$ and $id_{\mathcal{F}_n Y}$, where $B \in \mathcal{F}_n Y$ and $t \in [0, 1]$. Indeed, $(\mathcal{F}_n H_1)(B, 0) = H_1(B, 0) = (f \circ g)(B) = (\mathcal{F}_n f \circ \mathcal{F}_n g)(B)$ for all $B \in \mathcal{F}_n Y$, and $(\mathcal{F}_n H_1)(B, 1) = H_1(B, 1) = B = id_{\mathcal{F}_n Y}$. Similarly, the mapping $(\mathcal{F}_n H_2)(A, t) = H_2(A, t)$ is a homotopy between $\mathcal{F}_n g \circ \mathcal{F}_n f$ and $id_{\mathcal{F}_n X}$, where $A \in \mathcal{F}_n X$ and $t \in [0, 1]$. Indeed, $(\mathcal{F}_n H_2)(A, 0) = H_2(A, 0) = (g \circ f)(A) = (\mathcal{F}_n g \circ \mathcal{F}_n f)(A)$ for all $A \in \mathcal{F}_n X$, and $(\mathcal{F}_n H_2)(A, 1) = H_2(A, 1) = A = id_{\mathcal{F}_n X}$. It means that $\mathcal{F}_n X$ and $\mathcal{F}_n Y$ are homotopically equivalent. Proposition 1 is proved.

Corollary 2. If spaces X and Y are homotopically equivalent, then the spaces $\mathcal{F}_n X$ and $\mathcal{F}_n Y$ are also homotopically equivalent.

By a *covariant homotopy functor* we mean a functor \mathcal{F} in the category **Top** of topological spaces and their continuous mappings satisfying

(*) \mathcal{F} preserves homotopy, that is, if a mapping $H(x, t)$ is a homotopy between continuous mappings $f, g : X \rightarrow Y$, then $F(H(x, t))$ is also a homotopy between mappings $F(f), F(g) : F(X) \rightarrow F(Y)$.

Theorem 2. The functor of n -fold symmetric product \mathcal{F}_n is a covariant homotopy functor.

Proof. Now we will show that the functor \mathcal{F}_n satisfies the above three conditions.

(i) Let id_X be identity mapping in the topological space X . Then we have that $(\mathcal{F}_n id_X)(A) = id_X(A) = A$. It means that the mapping $\mathcal{F}_n id_X$ is an identity mapping in the topological space $\mathcal{F}_n X$.

(ii) Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be the continuous mappings. Then it follows that $\mathcal{F}_n(g \circ f)(A) = (g \circ f)(A) = g(f(A)) = (\mathcal{F}_n g)(f(A)) = \mathcal{F}_n g(A) \circ \mathcal{F}_n f(A)$.

(*) It follows easily from Theorem 1. Theorem 2 is proved.

REFERENCES

1. Beshimov R. B., Safarova D. T. Some Topological Properties of a Functor of Finite Degree. Lobachevskii of Journal Mathematics, 2021, V.42, P.2744–2753.
2. Beshimov R. B., Safarova D. T. Uniformly linked space and its hyperspace. AIP Conference Proceedings, 2024, V.3004(1), P.020003.
3. Kočinac Lj. D. R., Mukhamadiev F. G., Sadullaev A. K. On some properties of the superextension functor. Herald of Institute Mathematics of the National Academy of Sciences of the Kyrgyz Republic, 2022, V.2, P.5–18.
4. Kočinac Lj. D. R., Mukhamadiev F. G., Sadullaev A. K. On the superextension functor. AIP Conference Proceedings, 2023, V.2879(1), P.020004.
5. Kočinac Lj. D. R., Mukhamadiev F. G., Sadullaev A. K. Some Cardinal and Geometric Properties of the Space of Permutation Degree. Axioms, 2022, V.11(6), P.290.
6. Mukhamadiev F. G. Some homotopy properties of the space of complete linked systems. Selected papers of the 2023 International Conference on Topology and its Applications. In press.
7. Fedorchuk V. V., Filippov V. V. opology of Hyperspaces and Its Applications. Znanie: Moscow, Russia, 1989. (In Russian)
8. Shchepin E. V. Functors and uncountable powers of compacta. Math. Surveys. 1981, V.36:3, P.1–7.

9. Good C., Macias S. Symmetric products of generalized metric spaces. *Topology Appl.* 2016, V.206, P.93–114.
10. Tuyen L. Q., Tuyen O. V., Kočinac Lj. D. R. The Vietoris hyperspace $F(X)$ and certain generalized metric properties. *Hacettepe Journal of Mathematics and Statistics*, 2024, V.53(2), P.356–366.
11. Tang Z., Lin S., Lin F. Symmetric products and closed finite-to-one mappings. *Topology and its Applications*, 2018, V.234, P.6–45.
12. Peng L. X., Sun Y. A study on symmetric products of generalized metric spaces. *Topology and its Applications*, 2017, V.231, P.411–429.
13. Tuyen L. Q., Tuyen O. V. A note on the hyperspace of finite subsets. *Fasciculi Math.* 2021, V.65, P.67–73.
14. Yuldashev T. K., Mukhamadiev F. G. The local density and the local weak density in the space of permutation degree and in Hattori space. *Ural Mathematical Journal*, 2020, V.6(2), P.108–116.
15. Kosniowski C. *A First Course in Algebraic Topology*. Cambridge University Press, 1980.
16. Munkres J. *Topology*, Second ed. Pearson Education Limited, Edinburgh, 2014.

РЕЗЮМЕ

Ushbu maqolada biz X fazoning n -darajali simmetrik ko'paytma fazosining ba'zi gomotopik xossalarini o'rganamiz. Agar $f, g : X \rightarrow Y$ akslantirishlar o'zaro gomotop bo'lsa, u holda $\mathcal{F}_n f, \mathcal{F}_n g : \mathcal{F}_n X \rightarrow \mathcal{F}_n Y$ akslantirishlar ham o'zaro gomotop ekanligi isbotlangan. Bundan tashqari, n -darajali simmetrik ko'paytma funktori \mathcal{F}_n kovariant gomotopik funktor ekanligi ko'rsatilgan. Shu bilan birga, agar X va Y fazolar gomotopik ekvivalent bo'lsa, u holda $\mathcal{F}_n X$ va $\mathcal{F}_n Y$ fazolar ham gomotopik ekvivalent ekanligi isbotlangan.

Kalit so'zlar: Funktor, n -darajali simmetrik ko'paytma, gomotopiya, gomotopik ekvivalentlik.

РЕЗЮМЕ

В данной работе изучаются некоторые гомотопические свойства пространства n -кратного симметрического произведения пространства X . Доказывается, что если отображения $f, g : X \rightarrow Y$ гомотопны, то отображения $\mathcal{F}_n f, \mathcal{F}_n g : \mathcal{F}_n X \rightarrow \mathcal{F}_n Y$ также гомотопны. Также показано, что функтор n -кратного симметрического произведения \mathcal{F}_n является ковариантным гомотопическим функтором. Кроме того, доказано, что если пространства X и Y гомотопически эквивалентны, то пространства $\mathcal{F}_n X$ и $\mathcal{F}_n Y$ также гомотопически эквивалентны.

Ключевые слова: Функтор, n -кратное симметричное произведение, гомотопия, гомотопическая эквивалентность.